

Preface

The 7th Indonesia Japan Joint Scientific Symposium (IJSS 2016), the 24th CEReS International Symposium, the 4th Symposium on Microsatellite for Remote Sensing (SOMIRES), and the Symposium on Innovative Microwave Remote Sensing were held on the Nishi-Chiba campus of Chiba University, Japan, during 20-24 November 2016. These symposia focused on providing a forum to share and discuss recent issues and developments in various fields of science and technology.

IJSS has been held since 2004 and now it has become one of the successful regional symposia co-organized by two countries – Indonesia and Japan. In IJSS 2016, a total of 160 papers that have gone through review process are presented. In each of the 29 sessions, enthusiastic and stimulating discussions have led to the exchange of innovative ideas and advancement of the state of knowledge among students and researchers from universities in Japan, Indonesia, and other countries.

Generous supports for the conference were provided by Kabupaten Siak, Bank RiauKepri, Katoro, APPJ, Dental Support, Chiba Soy Sauce Union, and Kominato Railway. On behalf of the IJSS local organizing team, I would like to express my sincere gratitude to their supports, which greatly contributed to the participation of young scientists.

Finally, I would like to thank all the proceedings team members who have dedicated their effort and time to bring the manuscripts into the form of a book. This book will serve as a long-lasting credit to the achievements of IJSS 2016.

Professor Hiroaki Kuze, Dr. Sc.,

General Chairman of IJSS 2016

Committees

Advisory Committee:

Prof. Takeshi Tokuhisa, President of Chiba University, Japan
Prof. Muhammad Anis, Rector of Universitas Indonesia, Indonesia

General Chairman

Prof. Hiroaki Kuze

Executive Committee

Prof. Andi Eka Sakya, Head of Badan Meteorologi, Klimatologi, dan Geofisika, Indonesia
Prof. Herry Suhardiyanto, Rector of Institut Pertanian Bogor, Indonesia
Prof. Kadarsah Suryadi, Rector of Institut Teknologi Bandung, Indonesia
Prof. Yos Johan Utama, Rector of Universitas Diponegoro, Indonesia
Prof. Dwikorita Karnawati, Rector of Universitas Gadjah Mada, Indonesia
Prof. Dwia Aries Tina Pulubuhu, Rector of Universitas Hasanuddin, Indonesia
Prof. Muhammad Anis, Rector of Universitas Indonesia, Indonesia
Prof. Tri Hanggono Achmad, Rector of Universitas Padjadjaran, Indonesia
Prof. Ketut Suastika, Rector of Universitas Udayana, Indonesia
Prof. H. Detri Karya, Rector of Universitas Islam Riau, Indonesia
Prof. Ravik Karsidi, Rector of Universitas Sebelas Maret, Indonesia

Sister Universities (*)

Badan Meteorologi, Klimatologi, dan Geofisika, Indonesia
Institut Pertanian Bogor, Indonesia
Institut Teknologi Bandung, Indonesia
Universitas Diponegoro, Indonesia
Universitas Gadjah Mada, Indonesia
Universitas Hasanuddin, Indonesia
Universitas Indonesia, Indonesia
Universitas Padjadjaran, Indonesia
Universitas Udayana, Indonesia

* Base on alphabet queue

Organizing Committee

Prof. Hiroaki Kuze, Chiba University, Japan
Prof. Kenichi Sakakibara, Chiba University, Japan
Prof. Josaphat Tetuko Sri Sumantyo, Chiba University, Japan
Prof. Shogo Shimazu, Chiba University, Japan
Prof. Niikura Ryoko, Chiba University, Japan
Prof. M. Takagaki, Chiba University, Japan
Prof. Hiroshi Nakagawa, Chiba University, Japan
Prof. Hitoshi Asanuma, Chiba University, Japan
Prof. Ryutaro Tateishi, Chiba University, Japan
Prof. Koichi Ito, Chiba University, Japan
Prof. Kenzo Nonami, Chiba University, Japan
Prof. Katsumi Hattori, Chiba University, Japan
Prof. Gunawan Wibisono, Universitas Indonesia, Indonesia
Prof. Eko Tjipto Rahardjo, Universitas Indonesia, Indonesia
Prof. Danang Sri Hadmoko, Universitas Gadjah Mada, Indonesia
Prof. Arif Marfai, Universitas Gadjah Mada, Indonesia
Prof. Muhammad Asvial, Universitas Indonesia, Indonesia

Prof. Ketut Wikantika, Institut Teknologi Bandung, Indonesia
Prof. I Made Mahendra, Universitas Udayana, Indonesia
Prof. Darsono, Universitas Sebelas Maret, Indonesia
Prof. Edison Munaf, Universitas Andalas, Indonesia
Prof. Nelson Pomalingo, Universitas Muhammadiyah Gorontalo, Indonesia
Prof. Ohneda Osamu, University of Tsukuba, Japan
Prof. Isamu Nagano, Komatsu College, Japan
Assoc. Prof. Dodi Sudiana, Universitas Indonesia, Indonesia
Assoc. Prof. Chiharu Hongo, Chiba University, Japan
Assoc. Prof. Handarto, Universitas Padjadjaran, Indonesia
Assoc. Prof. Emma Hisbaron, Universitas Gadjah Mada, Indonesia
Assoc. Prof. Muhammad Isman Jusuf, Universitas Muhammadiyah Gorontalo, Indonesia

Technical Program Committee

Prof. Agus Hartoko, Universitas Diponegoro, Indonesia
Prof. Rika Haryana, Universitas Gadjah Mada, Indonesia
Prof. Andriyan Bayu Suksmono, Institut Teknologi Bandung, Indonesia
Prof. Wolfgang Martin Boerner, University of Illinois, United States
Prof. Keiichiro Kondo, Chiba University, Japan
Prof. Nobuo Takeuchi, Chiba University, Japan
Prof. Motoi Machida, Chiba University, Japan
Prof. Hideki Nakagome, Chiba University, Japan
Prof. Wuled Lenggoro, Tokyo Agricultural University, Japan
Prof. Bondan T. Sofyan, Universitas Indonesia, Indonesia
Prof. Sugeng, Universitas Islam Riau, Indonesia
Prof. Mukhtar Ahmad, Universitas Islam Riau, Indonesia
Prof. I Wayan Sandi Adnyana, Universitas Udayana, Indonesia
Prof. Purnomo Priambodo, Universitas Indonesia, Indonesia
Assoc. Prof. Kazuhiko Ohnuma, Chiba University, Japan
Assoc. Prof. Rie Ono, Chiba University, Japan
Assoc. Prof. Basari, Universitas Indonesia, Japan
Assoc. Prof. Hitoshi Irie, Chiba University, Japan
Assoc. Prof. Agfianto Putra, Universitas Gadjah Mada, Indonesia
Assoc. Prof. Yoshimasa Amano, Chiba University, Japan
Assoc. Prof. Kazuteru Namba, Chiba University, Japan
Assoc. Prof. Idris Mandang, Universitas Mulawarman, Indonesia
Assoc. Prof. Tri Kuntoro Priyambodo, Universitas Gadjah Mada, Indonesia
Assoc. Prof. Gede Karang, Universitas Udayana, Indonesia
Assoc. Prof. Masuda Kenji, Shizuoka University, Japan
Assoc. Prof. Katoh Kentaroh, Tsuruoka College, Japan
Assoc. Prof. Teti Zubaidah, Universitas Mataram, Indonesia
Assoc. Prof. Rahmatulloh, Universitas Indonesia, Indonesia
Assoc. Prof. Dr. Nurman, S.Si, M.Si, Universitas Islam Riau, Indonesia
Dr. Yohandri, Universitas Negeri Padang, Indonesia
Dr. Elyas Palantei, Universitas Hasanuddin, Indonesia
Dr. Ilham Alimuddin, Universitas Hasanuddin, Indonesia
Dr. Luhur Bayuaji, Universiti Malaysia Pahang, Malaysia
Dr. Muhammad Fauzan, Universitas Brawijaya, Indonesia
Dr. Yuhendra, Institut Teknologi Padang, Indonesia
Dr. Indra Riyanto, Universitas Budiluhur, Indonesia
Dr. Wahyudi Parnadi, Institut Teknologi Bandung, Indonesia
Dr. Robertus Heru, Lembaga Antariksa dan Penerbangan Nasional, Indonesia



The 7th Indonesia Japan
Joint Scientific Symposium
(IJSS 2016)
Chiba, 20-24 November 2016

Dr. Evizal Abdul Kadir, Universitas Islam Riau, Indonesia
Dr. Muhamad Komarudin, Universitas Lampung, Indonesia

**Local Steering Committee
Secretariat and Registration:**

Treasurer:
Chihaya Miyamoto

Publication:
Husnul Kausarian
Asif Awaluddin
Dr. Ratih Fitria
Putri Agus Hendra
Pakhrur Razi

Reviewers:
Prof. Kenichi Sakakibara
Prof. Rie Ono
Prof. Shogo Shimazu
Prof. Hitoshi Asanuma
Prof. Josaphat Tetuko Sri Sumantyo
Prof. Ryutaro Tateishi
Prof. Katsumi Hattori
Prof. Eko Tjipto Rahardjo
Prof. Danang Sri Hadmoko
Prof. Arif Marfai
Prof. Muhammad Asvial
Prof. Ketut Wikantika
Prof. I Made Mahendra
Prof. Agus Hartoko
Prof Hartono
Prof. Andriyan Bayu Suksmono
Prof. Motoi Machida
Prof. Kazuhiko Ohnuma
Prof. Kazuteru Namba
Prof. Idris Mandang
Prof. Gede Karang
Dr. Dodi Sudiana
Dr. Yohandri
Dr. Elyas Palantei
Dr. Ilham Alimuddin
Dr. Wahyudi Parnadi
Dr Mya Dwi Rostika
Dr Emmy Latifah

Website Coordinator:
Babag Purbantoro
Good Fried Panggabean
Farohaji Kurniawan

Documentation:
Heein Yang

Oral Session

Wednesday, November 23
09:30-15:00

09:30-12:00 Wed, Nov 23	Novel Disaster Mitigation Engineering Toward Future	Chair : Hiroshi Asanuma 3F Reception Hall
09:30 - 10:00 P132	Hiroshi Asanuma (p:1-9) <i>Smart disaster mitigation based on novel structures/materials</i>	
10:00 - 10:30 P055	Nanang T. Puspito (p:10-16) <i>Seismological Evidence for Crustal Deformation beneath the Sunda-Banda Arc Transition Zone</i>	
10:30 - 11:00 P126	Djati Mardiatno (p:17-25) <i>Identification of Area Prone to Multi Hazards using Remotely Sensed Data - Case in Lowland Area of Kulon Progo, Yogyakarta, Indonesia</i>	
11:00 - 11:30 P107	Toha Saleh (p:26-36) <i>Analysis of Stream and Residential Area Protection Case Study: Tembagapura, Papua</i>	
11:30 - 12:00 P057	Dyah Rahmawati Hizbaron (p:37-45) <i>Vulnerability of Volcanic Ternate Island Towards Ecosystem Based Disaster Risk Management</i>	
09:30-11:50 Wed, Nov 23	The 24th CEReS International Symposium	Chair : Nobuo TAKEUCHI 3F Meeting Room 4
09:30 - 09:50 P093	Nobuo TAKEUCHI (p:1-10) <i>Investigation of a High Resolution Lidar Using Optical Frequency Comb</i>	
09:50 - 10:10 P163	Nofel Lagrosas (p:11-14) <i>Modeling of scattering enhancement factor, $f(RH)$, in Chiba using visibility and ground measurements</i>	
10:10 - 10:30 P069	D. Ichikawa <i>Geostationary HIMAWARI-8 and Polar Orbiting S-NPP VIIRS Satellites for regional environmental monitoring of Indonesia</i>	
10:30 - 10:50 P041	Jamrud Aminuddin (p:15-18) <i>Observation of Aerosol Optical Properties by Means of Himawari-8 Satellite from Space and Lidar System from Surface</i>	
10:50 - 11:10 P110	Babag Purbantoro (p:19-22) <i>Cloud Retrieval and Cloud Type Detection from Himawari-8 Satellite Data Based on the Split Window Algorithm</i>	
11:10 - 11:30 P119	Ginaldi Ari N / Asif Awaludin (p:33-39) <i>Hail Detection By A Low Cost Local Weather Radar Operated For Disaster Early Warning System</i>	
11:30 - 11:50 P042	Andung Bayu Sekaranom (p:23-32) <i>Extreme precipitation over Indonesian maritime continent: Uncertainties in satellite estimation and its relationship with low storm top height extreme</i>	

Hail Detection By A Low Cost Local Weather Radar Operated For Disaster Early Warning System

Ginaldi Ari Nugroho¹, Tiin Sinatra¹, Asif Awaludin², Halimurrahman¹, Dian Handiana³

¹⁾ Center for Atmospheric Science and Technology, Lembaga Penerbangan dan Antariksa Nasional (LAPAN)

²⁾ Chiba University

³⁾ Indonesia Agency for Meteorology, Climatology, and Geophysics - BMKG

Email : ginaldi.ari@lapan.go.id

Abstract

A local weather radar (LWR) has been developed by modifying X-band marine radar. LWR has been installed and operated to measure precipitation in Bandung area, West Java, Indonesia at 107.586491° W and 6.894958° S. During its observation, LWR has detected hail falling in West Bandung region on March 26, 2016. LWR can observe hail formation and growth processes thanks to its 2 minutes of sampling interval. Hail occurrence was also confirmed by C-band weather radar installed 120 km away in northwest side of LWR site and black body temperature data of Himawari 8 satellite which analyzed using Auer method. This C-band radar identify hail occurrence at 2.43 pm until 3.14 pm (around 31 minutes). Maximum recorded reflectivity data of this hail precipitation is 58.46 dBZ which close to 55.1 dBZ of C-band radar data. This results confirm that LWR is suitable as a low cost solution for disaster early warning system due to precipitation.

Keywords

Keywords : local weather radar, Himawari 8, hail, reflectivity, disaster.

1. Introduction

Deep convective cloud mostly appear in tropical region. This convective cloud can generate severe weathers such as heavy rain, tornado, strong wind and hail. Urban areas are severe weather catastrophe-prone regions due to its dense population and building. Therefore, efforts to predict and monitor severe weather are necessary to avoid huge damage.

Hail, which formed by strong and wide updraft, has diameter until around 5 mm (Barnes, 2010). Many methods have been developed to detect hail. Weather Radar is a useful tool to conduct real time observation of hail. Dual polarization weather radar is very effective for distinguishing rain and hail by measuring Z_{DR} - differential reflectivity (Marzano et al., 2007). Single-polarization weather radars can not distinguish among different types of hydrometeors, however many method have been developed in order to distinguish between hail and rain using single polarization (Holleman, 2001). Most

single polarization radar is commonly S-Band or C-band system. Recently, there are low-cost short range X-Band systems developed for observing precipitation in urban area (Einfalt T, et al,2004 ; Rollenbeck and Bendix, 2006 ;Van de Beck et al., 2010) and hail storm (Capozzi, V et al, 2016).

In this paper, a preliminary results of using LWR and image processing for hail detection are presented. Hail with rain and strong wind has occurred in urban area of Cimahi city, a west region of Bandung in Indonesia, for around 30 minutes (14.45 pm) in March 26, 2016 (<http://jabar.tribunnews.com> , access May 20th 2016). The hail diameter was up to 2 cm with strong wind damaging trees and billboard. A local Weather Radar (LWR) developed in 2012 has been utilized to observe the precipitation around the radar site. The LWR is a precipitation radar developed by modifying an X-band marine radar. Signal and image processing system of the X-band radar are modified to analyze rainfall reflectivity data rather than marine target. Hail observation results were verified by comparing them with measurement results of a C-band weather radar located 120 km northwest of the LWR location.

2. Data and Methods

The LWR is a modification results of a marine radar. Data acquisition and digital signal processing are developed for the LWR to extract, overlay, and display precipitation reflectivity (Nugroho and Awaludin, 2013). In order to verify the hail occurrence, observation data from Himawari 8 satellite and Indonesia Agency for Meteorology, Climatology, and Geophysics (BMKG) C-Band weather Radar in March 26, 2016 are utilized. The C-Band weather radar has single polarization and its coverage area compared to the LWR area is illustrated in Fig. 1. White dot is the location of the C-Band radar while the yellow pindrop is the LWR location.



Figure 1. Coverage area of C-Band Radar (green circle) and LWR (red circle).

C-band weather radar with CAPPI method and black body temperature (T_{bb}) data from Himawari satellite were used to identify the hail object. Previous research showed that increase on diameter of scattering particles will increase radar reflectivity significantly. Threshold was used to distinguish between severe rain and hail by employing CAPPI method with 57 dBZ (Mason 1971) and 54 dBZ (Auer,1994) of threshold. Obtained hail properties were compared with LWR observation data. Location and reflectivity value were analyzed to know the LWR performance in detecting hail.

3. Results and Discussions

A method to recognize hail by using combination of radar reflectivity and cloud-top temperature had been proposed in (Auer, 1994). According to this method, threshold of CAPPI reflectivity (Z_{TH}) for hail as a function of cloud-top temperature (T_{top}) is given by equation (1). Optimum threshold for this equation varies between 36 and 53 dBZ for cloud-top temperatures between -11 and -55°C .

$$Z_{TH} = \begin{cases} -0.38.(T_{top} - 85.0) & \text{if } T_{top} \leq -11^{\circ}\text{C} \\ 1.33.(T_{top} + 38.8) & \text{if } T_{top} > -11^{\circ}\text{C} \end{cases} \quad (1)$$

Hail occurrence in Cimahi city, west region of Bandung, was investigated using this method. Black body temperature (T_{bb}) data of Himawari 8 satellite was presented as cloud-top temperature data as shown in Fig. 2. This data reveals that cloud with top temperature around -60°C appears during hail occurrence which indicates strong convective process. Z_{TH} value of this temperature is 55.1 dBZ.




Figure 2. Temperature Black Body (T_{bb}) data of Himawari-8 satellite at 08:00 UTC.





Figure 3. Reflectivity in CAPPI by C-Band weather radar at 4.25 dan 4.75 km.

Observation data by C-band weather radar at constant altitude of 4.25 and 4.75 km were presented to confirm Himawari 8 data. Both radar and satellite data confirm high reflectivity above 50 dBZ at 08:00 UTC but with slightly different location of west Bandung region. Whereas colocation data of both radar and satellite reveals reflectivity

threshold of 40.77-46.38 dBZ in that region. Since cloud top temperature in the collocation area is below -30°C , it can be concluded as hail occurrence.






Figure 4. Precipitation evolution from the LWR observation data.

C-Band weather radar data indicated that high reflectivity is detected in west Bandung region. LWR also observed high reflectivity in similar location. Time frame images of both radar showed similar time period of high reflectivity between 07.40 until 08.10 UTC. Thus, according to both radars, hail occurred for about 1 hour 40 minutes. However, the shape of the reflectivity area from both radar are different, this is due to different specification of both radar and also scan method. By using time sequential images as outlined in Fig. 4, it can be observed that the growth of the heavy precipitation (including hail) started at 07.18 UTC, continued to grow at 08.04 UTC, and then dispersed at 08.58 UTC.

Observed LWR reflectivity data is much lower than C-band weather radar results for similar target due to its low power and wide beamwidth. In this regard, 4 kW LWR measurement data has been calibrated using 25 kW X-band weather radar (Nugroho, 2015). According to the calibration results, LWR reflectivity which equal to the X-band radar reflectivity is given by

$$dBZ_{LWR} = 7.338 \times \ln(ADC \text{ count}) + 22.71 \quad (2)$$

By using equation (2), calibration result of 30 dBZ reflectivity in LWR data which has 4,8721 of ADC count is 58.46 dBZ, slightly higher than 55.1 dBZ of C-band radar reflectivity. Both radar data confirm hail occurrence since minimum reflectivity that confirm hail by radar is 50 dBZ (Auer, 1994).

4. Conclusion

Hail occurrence during heavy precipitation in March 26, 2016 has been investigated using developed LWR. Black body temperature (T_{bb}) data of Himawari 8 satellite and images data of C-band weather radar data are presented to verify hail detection of the

LWR results using Auer method. Both satellite and C-band radar data confirm the hail occurrence. LWR observation result is in good agreement with C-band radar data in hail detection by recording LWR maximum reflectivity of 58.46 dBZ compared to 55.1 dBZ of C-band radar data.

Acknowledgements

This research is supported by Center for Atmospheric Science and Technology, Lembaga Penerbangan dan Antariksa Nasional (LAPAN) Indonesia for the LWR project and Indonesia Agency for Meteorology, Climatology, and Geophysics - BMKG for C-Band weather radar data.

References

- Barnes, G. M., (2010) Meteorological hazards in the Tropics: Severe convective storms and flash floods. Chapter in Tropical Meteorology, Encyclopedia of Life Support Systems (EOLSS) (www.eolss.net), sponsored by the UNESCO, 109 pp.
- Nugroho, G.A., Munir, M.M., dan Khairurrujal. (2015) A Computer-Based Marine Automatic Radar for Rain detection. *Applied Mechanics and Materials*, **771**, 9-12
- Awaludin, A., G.A. Nugroho, dan Rahayu, S.A. (2013) Analisis Kemampuan Radar Navigasi Laut Furuno 1932 Mark-2 untuk Pemantauan Intensitas Hujan. *Jurnal Sains Dirgantara*, **10**, 90-103.
- Auer, A. H., Jr: 1994, Hail recognition through the combined use of radar reflectivity and cloud-top temperatures. *Mon. Wea. Rev.*, **122**, 2218-2221.
- Mason, B. J.: 1971, *The Physics of Clouds*. Clarendon Press, Oxford UK.
- Iwan Holleman Hail detection using single-polarization radar. 2001. Scientific Report, KNMI WR-2001-01.
- Nugroho, G A and Awaludin A. (2013). Mapping method development using digital image processing to calibrate rainfall radar image, International Proceeding 17th SIPTEKGAN.
- Marzano, F.S., Budillon, G., Picciotti, E., Montopoli, M., Zinzi, A., and Buonocore, B.: X-band weather radar monitoring realtime products in Rome and Naples urban areas. *Tyrrhenian Workshop 2012 on Advances in Radar and Remote Sensing, 2012*
- Marzano F.S., Scaranari, D., and Vulpiani, G.: Supervised fuzzy-logic classification of hydrometeors using C-band dualpolarized radars. *IEEE Trans. Geosci. Rem. Sensing*, ISSN: 0196-2892, n. 45, pp. 3784-3799, 2007
- Einfalt T, et al, (2004), Towards a roadmap for use of radar rainfall data in urban drainage, *Journal of Hydrology*, **299**, 186-202.
- Rollenbeck, R. dan Bendix, J., (2006), Experimental calibration of a cost-effective X-band weather radar for climate ecological use in southern Ecuador, *Atmospheric Research*, **79**, 296-316
- Van de Beek, C. Z., Leijnse, H., Stricker, J. N. M., Uijlenhoet, R., and Russchenberg, H. W. J.: Performance of high-resolution X-band radar for rainfall measurement in The Netherlands. *Hydrol. Earth Syst. Sci.*, Vol. 14, pp. 205-221, 2010
- Vincenzo Capozzi, Errico Picciotti, Vincenzo Mazzarella, Giorgio Budillon, Frank Silvio Marzano, *journal Hydrol. Earth Syst. Sci.*, doi:10.5194/hess-2016-177, 2016
<http://jabar.tribunnews.com>, access May 20th 2016